

A short note on $q\bar{q}g$ hybrid assignment for $X(1812) \rightarrow \omega\phi$

Kuang-Ta Chao^(a,b)

(a) Department of Physics, Peking University, Beijing 100871, People's Republic of China

(b) China Center of Advanced Science and Technology (World Laboratory), Beijing 100080, People's Republic of China

Abstract

BES Collaboration has recently observed an enhancement near the $\omega\phi$ threshold in the double OZI suppressed decay $J/\psi \rightarrow \gamma\omega\phi$. This $\omega\phi$ enhancement is probably due to a new $J^{PC} = 0^{++}$ $X(1812)$ state. We discuss a possible assignment that the $X(1812)$ is a $1/\sqrt{2}(u\bar{u} + d\bar{d})g$ (or $s\bar{s}g$) hybrid meson. We suggest further measurements to test these assignments.

Based on a sample of 5.8×10^7 J/ψ events, very recently the BES Collaboration has observed an enhancement near threshold in the $\omega\phi$ invariant mass spectrum from the double OZI suppressed decay of $J/\psi \rightarrow \gamma\omega\phi$ with a statistical significance of more than 10σ . As a hadronic resonance state X , this enhancement has the following mass, width, and branching ratio product[1]:

$$M = 1812_{-26}^{+19} \pm 18 \text{ MeV}, \quad \Gamma = 105 \pm 20 \pm 28 \text{ MeV}, \quad (1)$$

$$B(J/\psi \rightarrow \gamma X) \times B(X \rightarrow \omega\phi) = (2.61 \pm 0.27 \pm 0.65) 10^{-4}. \quad (2)$$

Moreover, the spin-parity $J^{PC} = 0^{++}$ is favored for $X(1812)$ by a partial wave analysis.

J/ψ radiative decays into light hadrons proceed through $J/\psi \rightarrow \gamma gg$ and are viewed as the glue-rich channels. The striking feature of the $X(1812)$ that it decays substantially to an ω , which is a $(1/\sqrt{2})(u\bar{u} + d\bar{d})$ meson, and a ϕ , which is a $s\bar{s}$ meson, may imply that the $X(1812)$ is unlikely to be an ordinary $q\bar{q}$ meson, because the $q\bar{q} \rightarrow q\bar{q} + q'\bar{q}'$ decays (q and q' are of different flavors) should be suppressed, compared with the OZI allowed $q\bar{q} \rightarrow q\bar{q}' + q'\bar{q}$ decays. Therefore, the $X(1812)$ is likely to be a new type of hadrons other than the conventional $q\bar{q}$ mesons.

In the following we will discuss a possible assignment for the $X(1812)$. That is, the $X(1812)$ is a $q\bar{q}g$ hybrid state.

The expected existence of hybrid states is an essential feature of quantum chromodynamics. A hybrid state $q\bar{q}g$ is made of $(q\bar{q})_8$, a quark and an antiquark in color-octet, combined with an excited gluon g . Hybrid states have been extensively studied in various approaches such as the constituent gluon model[2], the flux tube model[3], the bag model[4, 5], the QCD sum rules[6], and the lattice QCD[7, 8]. The experimental status for hybrid states can be found in a mini-review for "Non- $q\bar{q}$ candidates" written by the Particle Data Group[9]. As a more fundamental theoretical approach for hybrid studies, lattice QCD has made interesting progress and shows that for an exotic spin-parity $J^{PC} = 1^{-+}$ hybrid, which does not mix with ordinary $q\bar{q}$ mesons

and is then a "pure" hybrid state, the mass is estimated to be in the range 2.0(2) GeV for s quarks, and 1.9(2) GeV for u, d quarks. The decay has also been studied for the heavy quark hybrid, for which the dominant decay mode is the string de-excitation into a flavor singlet meson, and the magnitude of the decay width is found to be of order 100 MeV[7]. Although the hadronic decays for light-quark hybrid mesons has not been explored, the above lattice QCD results do shed light on a rough estimate of decay widths and modes for light-quark hybrids. Concerning the X(1812) observed by BES, it seems that its mass, total width, and decay modes in particular, might be compatible with a 0^{++} light-quark $(1/\sqrt{2})(u\bar{u} + d\bar{d})g$ hybrid state. In the flux tube model the mass of 0^{++} hybrid is higher than that of 1^{-+} ($1^{+-}, 1^{++}, 1^{--}, 0^{+-}, 0^{-+}, 2^{+-}, 2^{-+}$ as well), but considering the exiting theoretical uncertainties including the effects of mixing with other 0^{++} states, a mass of 1812 MeV for the 0^{++} hybrid does not seem to be unacceptable. At present it is still premature to make precise quantitative predictions for hybrid states due to our limited understanding of nonperturbative dynamics. Therefore we will focus on some distinctively qualitative features of the hybrid states and discuss a possible hybrid meson interpretation for the X(1812).

As already noted in [5, 12], the decay into $\omega\phi$ can be a distinctive feature of 0^{++} and 2^{++} $q\bar{q}g$ hybrid states including both $(1/\sqrt{2})(u\bar{u} + d\bar{d})g$ [5] and $s\bar{s}g$ states[12]. Since for the X(1812) the $\omega\phi$ mode is the only decay channel observed so far, it is natural to propose that the X(1812) be a possible candidate of the 0^{++} $q\bar{q}g$ states.

We first use a bag model estimation. In the spherical approximation of the bag model, the $q\bar{q}$ can couple to either the TE (transverse electric) or the TM (transverse magnetic) gluon. The TE mode has an axial vector quantum number $J^{PC} = 1^{+-}$, while the TM mode has $J^{PC} = 1^{-+}$ [5]. Moreover, in this model it was found that the TE gluon couples in the s-channel to $u\bar{u}, d\bar{d}$, and $s\bar{s}$ in an approximately flavor symmetric way, whereas the TM gluon couples to $s\bar{s}$ much stronger than to $u\bar{u}$ and $d\bar{d}$ [5]. In this model, we may assign X(1812) as the 0^{++} $(1/\sqrt{2})(u\bar{u} + d\bar{d})g_{TM}$ state, which has a mass in the range 1.51-1.90 GeV. Note that in this model the gluon excitation g_{TM} couples dominantly to a color-octet $s\bar{s}$, so we would approximately have

$$g_{TM} \rightarrow (s\bar{s})_8, \quad (3)$$

and expect this hybrid to decay via

$$X(1812) = \frac{1}{\sqrt{2}}(u\bar{u} + d\bar{d})g_{TM} \rightarrow \omega_8\phi_8 \rightarrow \omega\phi, K^*\bar{K}^*, K\bar{K}, \dots \quad (4)$$

Here the color-octet quark pairs $\omega_8\phi_8$ hadronize into color-singlet meson pairs by gluon exchange into $\omega\phi$ (without spin flip), and $\eta\eta, \eta\eta'$ (with spin flip) or quark rearrangement into $K^*\bar{K}^*, K\bar{K}$. The long-ranged color-electric force can induce the non-spin-flip transition, whereas short-ranged color-magnetic force causes the spin-flip transition. If the long-ranged color-electric force is stronger than the short-ranged color-magnetic force, we would expect the $\omega\phi$ decay to be stronger than the $\eta\eta, \eta\eta'$ decays. Furthermore, if the hybrid decay proceeds mainly via string de-excitation into a flavor singlet meson, as indicated by the lattice calculation for the heavy quark hybrid meson (the 1^{-+} charmonium hybrid $c\bar{c}g$ mainly dceays into $\chi_{c1} + \sigma$ with $\sigma \rightarrow \pi\pi$)[7], we would expect the $\omega\phi$ decay to be also stronger than the $K^*\bar{K}^*, K\bar{K}$ decays. One of the distinctive features in this model is that the decay into $\omega\omega$ will be much weaker than into $\omega\phi$. Moreover, in this model, we would predict the existence of the s quark partner of the u, d quark hybrid X' , and its decay modes

$$X' = s\bar{s}g_{TM} \rightarrow \phi_8\phi_8 \rightarrow \phi\phi, \eta\eta, \eta\eta', \eta'\eta', \dots \quad (5)$$

Here the X' has $J^{PC} = 0^{++}$ and its mass may be roughly estimated to be ranging from 2000 to 2100 MeV by taking the mass difference between s quark and u, d quark to be 100-150 MeV.

However, if the mass of X' is below the $\phi\phi$ threshold, its $\phi\phi$ decay mode would not be observed. In any case, if this model prediction makes sense, a weak signal of $0^{++} X(1812) \rightarrow \omega\omega$, and a strong signal of $0^{++} X'(2040 - 2100) \rightarrow \phi\phi$ should be observable experimentally. We note that the MarIII Collaboration[10] reported that the $J/\psi \rightarrow \gamma\omega\omega$ decay has a branching ratio of $(1.76 \pm 0.09 \pm 0.45) \times 10^{-3}$ and the $\omega\omega$ invariant mass distribution peaks at 1.8 GeV, but the $\omega\omega$ system below 2 GeV is predominantly pseudoscalar. Similar results were also reported by the DM2 Collaboration[11]. It is therefore very desirable to check the BES data for this channel to see if there is a $J^{PC} = 0^{++} \omega\omega$ component in the peak of 1.8 GeV.

We now turn into a more general discussion by assuming that the gluon couples to the color-octet $u\bar{u}, d\bar{d}, s\bar{s}$ symmetrically in the decay of hybrid state $(1/\sqrt{2})(u\bar{u} + d\bar{d})g$

$$g \rightarrow 1/\sqrt{3}(u\bar{u} + d\bar{d} + s\bar{s})_8. \quad (6)$$

If this is the case, then the $X(1812)$ would have stronger decay into $\omega\omega$ than $\phi\phi$, and stronger decay into $\pi\pi$ than $K\bar{K}$. These predictions may also be tested by experiment.

We might also interpret the $X(1812)$ as the $s\bar{s}g$ hybrid meson. Then it would not decay into $\omega\omega$. However, its mass is too small, compared with most predictions of theoretical models and lattice QCD calculations.

Note added. Related to the BES result recently reported in [1], a paper[13] proposed a four-quark state interpretation for the $X(1812)$; and another subsequent paper[14] focused on the mixing between $0^{++} q\bar{q}$ mesons, glueballs, and $q\bar{q}g$ hybrids, and used the $f_0(1790)=1/\sqrt{2}(u\bar{u} + d\bar{d})g$ as input and then found $X(1812)$ to be mainly the $s\bar{s}g$ hybrid state.

Acknowledgments. The author would like to thank X.Y. Shen for valuable discussions on the feasibility for experimental test of the proposed hybrid states. This work was supported in part by the National Natural Science Foundation of China (No 10421503), the Key Grant Project of Chinese Ministry of Education (No 305001), and the Research Found for Doctorial Program of Higher Education of China.

References

- [1] M. Ablikim et al. (BES Collaboration), hep-ex/0602031.
- [2] D. Horn and J. Mandula, Phys. Rev. D17, 898 (1978).
- [3] N. Isgur and J. Paton, Phys. Rev. D31, 2910 (1985); N. Isgur, R. Kokoski, and J. Paton, Phys. Rev. Lett. 54, 869 (1985);
F.E. Close and P.R. Page, Nucl. Phys. B443, 233 (1995); T. Barnes, F.E. Close, and E.S. Swanson, Phys. Rev. D52, 5242 (1995); F.E. Close and S. Godfrey, Phys. Lett. B574, 210(2003);
- [4] T. Barnes and F.E. Close, Phys. Lett. B116, 365 (1982).
- [5] M.S. Chanowitz and S.R. Sharpe, Phys. Lett. B132, 413 (1983); Nucl. Phys. B222, 211 (1983).
- [6] See, e.g., J. Govaerts et al., Nucl. Phys. B248, 1 (1984); F. de Viron and J. Govaerts, Phys. Rev. Lett. 53, 2207 (1984).
S.L. Zhu, Phys. Rev. D60, 014008 (1999); D60, 097502 (1999).

- [7] For a review on the lattice QCD study of hybrid states, see C. Michael, hep-ph/0308293, and references therein.
- [8] For some recent studies, see also, X.Q. Luo and Y. Liu, hep-lat/0512044; T.W. Chiu and T.H. Hsieh, hep-lat/0512029.
- [9] S. Eidelman et al., Phys. Lett. B592, 1 (2004).
- [10] R.M. Baltrusaitis et al., Phys. Rev. Lett. 55, 1723 (1985).
- [11] D. Bisello et al., Phys. Lett. B192, 239 (1987).
- [12] K.T. Chao, Phys. Rev. Lett. 60, 2579 (1988).
- [13] B.A. Li, hep-ph/0602072.
- [14] X.G. He, X.Q. Li, X. Liu, and X.Q. Zheng, hep-ph/0602075.